

**MEDIQUE AT HOME IPRIN- ibuprofen tablet, film coated**  
**Unifirst First Aid Corporation**

-----  
**Medique @ Home Iprin Tablets**

***Drug Facts***

***Active ingredient (in each tablet)***

Ibuprofen 200 mg (NSAID\*)

\*nonsteroidal anti-inflammatory drug

***Purpose***

Pain reliever/fever reducer

***Uses***

temporarily relieves minor aches and pains associated with:

- headache ■ toothache ■ backache ■ menstrual cramps
- common cold ■ muscular aches ■ minor arthritis pain

temporarily reduces fever

***Warnings***

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives ■ skin reddening ■ asthma (wheezing) ■ facial swelling ■ rash ■ shock ■ blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning :** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Do not use**

- if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever

reducer

- right before or after heart surgery

**Ask a doctor before use if**

- you have problems or serious side effects from taking pain relievers or fever reducers
- stomach bleeding warning applies to you
- you have a history of stomach problems such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma or had a stroke
- you are taking a diuretic

**Ask a doctor or pharmacist before use if you are**

- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- under a doctor's care for any serious condition
- taking any other drug

**When using this product**

- take with food or milk if stomach upset occurs

**Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- you have symptoms of heart problems or stroke
  - chest pain
  - trouble breathing
  - weakness in one part or side of body
  - slurred speech
  - leg swelling
- pain gets worse or lasts for more than 10 days
- fever gets worse or lasts for more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless specifically directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Directions**

- do not use more than directed

■ **the smallest effective dose should be used**

■ do not take longer than 10 days, unless directed by a doctor (see Warnings)

**Adults and children: (12 years and older)**

■ take 1 tablet every 4 to 6 hours while symptoms persist

■ if pain or fever does not respond to 1 tablet, 2 tablets may be used

■ do not exceed 6 tablets in 24 hours, unless directed by a doctor.

**Children under 12 years:**

■ ask a doctor

***Other information***

■ store at 68-77°F (20-25°C)

■ avoid excessive heat 104°F (above 40°C)

■ **TAMPER EVIDENT** child resistant packets

■ do not use any opened or torn packets

***Inactive ingredients***

carnauba wax\*, corn starch, hypromellose\*, iron oxide red, lactose\*, magnesium stearate\*, microcrystalline cellulose\*, polydextrose\*, polyethylene glycol, polyvinyl alcohol\*, povidone k-30\*, silicon dioxide, sodium starch glycolate, stearic acid, talc\*, titanium dioxide

\*may contain

***Questions or comments?*** 1-800-634-7680

Medique® @ Home

I-Prin Tablets

Ibuprofen 200 mg

Pain Reliever/Fever Reducer

(NSAID)

Pull to Open

Child-Resistant Packets

\*not actual size

50 Packets

(2 Tablets per Packet)

Compare to the active ingredient in Advil® Tablets

**Medique® @ Home**  
**I-Prin Tablets**  
Ibuprofen 200 mg  
Pain Reliever/Fever Reducer  
(NSAID)

Child-Resistant Packets  
\*not actual size

Compare to the active ingredient in Advil® Tablets

**The Brands That Work™ Now Work At Home**

100 Tablets 50 Packets (2 Tablets Per Packet)

**Medique® @ Home**  
**I-Prin Tablets**  
Ibuprofen 200 mg  
Pain Reliever/Fever Reducer  
(NSAID)

Child-Resistant Packets  
\*not actual size

Compare to the active ingredient in Advil® Tablets

50 Packets (2 Tablets Per Packet)

**Drug Facts**

**Active ingredient (in each tablet) Purpose**  
Ibuprofen 200 mg (NSAID) Pain reliever/fever reducer  
Nonsteroidal anti-inflammatory drug

**Uses**  
Temporarily relieves minor aches and pains associated with:  
• Headache • Toothache • Backache • Menstrual cramps  
• Common cold • Muscular aches • Minor arthritis pain  
Temporarily reduces fever

**Warnings**  
Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:  
• Hives • Skin redness/itching • Swelling (face/throat)  
• Facial swelling • Rash • Shock • Stomach  
If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:  
• Are age 65 or older  
• Have had stomach ulcers or bleeding problems  
• Take a blood thinner (anticoagulant) or steroid drug  
• Take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)  
• Have 3 or more alcoholic drinks every day while using this product  
• Use more or for a longer time than directed

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Do not use**  
• If you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer  
• Right before or after heart surgery

**Drug Facts (continued)**

**Ask a doctor before use if**  
• You have problems or serious side effects from taking pain relievers or fever reducers  
• Stomach bleeding warning applies to you  
• You have a history of stomach problems such as heartburn  
• You have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma or had a stroke  
• You are taking a diuretic

**Ask a doctor or pharmacist before use if you are**  
• Taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin  
• Under a doctor's care for any serious condition  
• Taking any other drug

**When using this product**  
• Take with food or milk if stomach upset occurs  
• Stop use and ask a doctor if:  
• You experience any of the following signs of stomach bleeding:  
• Stomach pain • Vomiting blood  
• Have bloody or black stools  
• Have stomach pain that does not get better  
• You have symptoms of heart problems or stroke  
• Chest pain • Trouble breathing  
• Weakness in one part or side of body  
• Pain gets worse or lasts for more than 10 days  
• Fever gets worse or lasts for more than 3 days  
• Redness or swelling is present in the painful area  
• Any new symptoms occur

**If pregnant or breast feeding:** Ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless specifically directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222) (continued on back panel)

**Drug Facts (continued)**

**Directions**  
• Do not use more than directed  
• The smallest effective dose should be used  
• Do not take longer than 10 days, unless directed by a doctor (see Warnings)

**Adults and children (12 years and older)**  
• Take 1 tablet every 4 to 6 hours while symptoms persist  
• If pain or fever does not respond to 1 tablet, 2 tablets may be used  
• Do not exceed 6 tablets in 24 hours, unless directed by a doctor

**Children under 12 years:**  
• Ask a doctor

**Other information**  
• Store at 68°F to 77°F (20° to 25°C)  
• Avoid excessive heat 104°F (above 40°C)  
• TAMPER EVIDENT Child-resistant packets  
• Do not use any opened or torn packets

**Inactive ingredients**  
Carbamide wax, corn starch, hydroxypropylcellulose, iron oxide red, lactose, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, polyvinyl alcohol, polyvinylpyrrolidone, silicon dioxide, sodium starch glycolate, stearic acid, talc, titanium dioxide

**Questions or comments?** 1-800-634-7680

1Advil® is a registered trademark of Pfizer Consumer and is not affiliated with Medique Products or this product.

Reorder 70833  
Manufactured for  
Medique Products  
Fort Myers, FL 33907  
1-800-634-7680  
www.mediqueproducts.com

Retain carton for complete product information

Medique® @ Home

I-Prin Tablets

Ibuprofen 200 mg

Pain Reliever/Fever Reducer

(NSAID)

Pull to Open

Child-Resistant Packets

\*not actual size

50 Packets

(2 Tablets per Packet)

Compare to the active ingredient in Advil® Tablets



STARCH, CORN (UNII: O8232NY3SJ)

Product Characteristics

Color	brown	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	44;291
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:47682-683-33	50 in 1 BOX	02/01/2021	
1		2 in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA075010	02/01/2021	

<b>MEDIQUE AT HOME IPRIN</b>			
ibuprofen tablet, film coated			
<b>Product Information</b>			
<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:47682-783
<b>Route of Administration</b>	ORAL		
<b>Active Ingredient/Active Moiety</b>			
<b>Ingredient Name</b>		<b>Basis of Strength</b>	<b>Strength</b>
<b>IBUPROFEN</b> (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)		IBUPROFEN	200 mg
<b>Inactive Ingredients</b>			
<b>Ingredient Name</b>			<b>Strength</b>
<b>POVIDONE K30</b> (UNII: U725QWY32X)			
<b>POLYVINYL ALCOHOL, UNSPECIFIED</b> (UNII: 532B59J990)			
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)			
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)			
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)			
<b>FERRIC OXIDE RED</b> (UNII: 1K09F3G675)			
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)			
<b>TALC</b> (UNII: 7SEV7J4R1U)			

<b>SODIUM STARCH GLYCOLATE TYPE A POTATO</b> (UNII: 5856J3G2A2)				
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)				
<b>Product Characteristics</b>				
<b>Color</b>	brown (Reddish Brown)	<b>Score</b>	no score	
<b>Shape</b>	ROUND	<b>Size</b>	10mm	
<b>Flavor</b>		<b>Imprint Code</b>	G;2	
<b>Contains</b>				
<b>Packaging</b>				
<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:47682-783-33	50 in 1 BOX	02/01/2021	
1		2 in 1 PACKET; Type 0: Not a Combination Product		
<b>Marketing Information</b>				
<b>Marketing Category</b>		<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
ANDA		ANDA079174	02/01/2021	

**Labeler** - Unifirst First Aid Corporation (832947092)

Revised: 5/2022

Unifirst First Aid Corporation